



TI 2006-022/1

Tinbergen Institute Discussion Paper

A Study on Linear Inequality Representation of Social Welfare Functions

Keisuke Sato

Yoshitsugu Yamamoto

University of Tsukuba, Japan, and Tinbergen Institute.

A STUDY ON LINEAR INEQUALITY REPRESENTATION OF SOCIAL WELFARE FUNCTIONS

KEISUKE SATO AND YOSHITSUGU YAMAMOTO

ABSTRACT. This paper presents a study on the recently proposed linear inequality representation of Arrovian Social Welfare Functions (ASWFs). We first give an alternative proof of the ASWF integer linear inequality representation theorem, and then show several sufficient conditions on preference domains for the linear inequalities of the representation to form integral polytopes. We also show that a given probabilistic ASWF induces a real vector satisfying the inequalities.

1. INTRODUCTION

Social choice theory discusses the way of aggregating individual opinions and making a decision of the society. Since Arrow [2] proved the classic Impossibility Theorem, this theory has been studied by various researchers of various academic fields. For the comprehension and the development of the theory see Arrow, Sen, and Suzumura [3] as well as Sen [10]. Recently Sethuraman, Teo, and Vohra [11] formulated Arrovian Social Welfare Functions (ASWFs), which are social welfare functions satisfying fundamental Pareto principle and independence of irrelevant alternatives axioms, as integer solutions to certain linear inequalities. They showed the Impossibility Theorem and other results such as Kalai and Muller [6] by solving their inequality system.

In this paper we study the linear inequality representation of ASWFs. We give an alternative proof of the one-to-one correspondence result between an ASWF and an integer solution to the ASWF linear inequality representation in Sethuraman et al. [11]. Next we consider some polyhedral structure determined by the linear inequalities of the representation. We correct one claim in Sethuraman et al. [11] and show that when the preference domain is single-peaked, single-caved, or the domain on which each triple of alternatives contains an alternative that cannot be medium, and it satisfies what we define as the “weakly nonisolated” condition, the set of nonnegative solutions to the inequalities forms an integral polytope. We then discuss noninteger solutions to the inequality system. After showing that a real vector satisfying the ASWF inequalities can be constructed from every “probabilistic” ASWF, we derive a subadditive function of Barberá and Sonnenschein [4] as a special case. The construction of a probabilistic ASWF from a given solution to the inequalities is also studied.

The rest of the paper is organized as follows. Section 2 introduces notations and axioms. In Section 3 we give an alternative proof of the ASWF integer linear inequality representation theorem of Sethuraman et al. [11]. Discussions as to some polyhedral structure determined by the linear inequalities are held in Section 4. In Section 5 we discuss a relation between probabilistic ASWFs and noninteger solutions to the linear inequality representation. Section 6 concludes the paper and we give our view to future work.

Date: February 20, 2006.

2. NOTATIONS AND AXIOMS

The finite set of *players* is denoted by $\mathcal{N} = \{1, 2, \dots, n\}$ and we suppose that the number of players is at least two. Let $\mathcal{X} = \{x, y, z, \dots\}$ be the set of all *alternatives* and assume $3 \leq |\mathcal{X}| < \infty$. A binary relation \succ on \mathcal{X} is a *linear ordering* if it satisfies

- (i) *completeness*: $x \succ y$, $y \succ x$, or both hold for any pair of alternatives $x, y \in \mathcal{X}$,
- (ii) *transitivity*: if $x \succ y$ and $y \succ z$ then $x \succ z$ holds for any alternatives $x, y, z \in \mathcal{X}$, and
- (iii) *antisymmetry*: if $x \succ y$ and $y \succ x$ then $x = y$ holds for any pair of alternatives $x, y \in \mathcal{X}$.

Let $L(\mathcal{X})$ be the family of all linear orderings on \mathcal{X} . The statement $x \succ y$ is read “ x is preferred to y .” We call a member of $L(\mathcal{X})$ a *preference ordering* or simply *preference*. For a subset $\mathcal{Y} \subseteq \mathcal{X}$, we denote by $\succ|_{\mathcal{Y}}$ the restriction of binary relation \succ to \mathcal{Y} , i.e., $\succ|_{\mathcal{Y}}$ is defined on $\mathcal{Y} \times \mathcal{Y}$ and $x \succ|_{\mathcal{Y}} y$ if and only if $x \succ y$ and $x, y \in \mathcal{Y}$.

Let Ω be a nonempty subset of $L(\mathcal{X})$ and call it the *preference domain* or the *domain*. We set $\mathcal{P} = \Omega^n$, the n -ary Cartesian product of Ω , call an element $p \in \mathcal{P}$ a *profile*, and denote by \succ_i^p the *preference of player i at profile p* . We denote by $\mathcal{S}(p, x \succ y)$ the set of players preferring alternative x to y at $p \in \mathcal{P}$, i.e.,

$$\mathcal{S}(p, x \succ y) := \{i \in \mathcal{N} \mid x \succ_i^p y\}.$$

Given $\mathcal{P} \subseteq L(\mathcal{X})^n$ we let $\mathcal{C}^{\mathcal{P}}(S, x \succ y)$ be the set of profiles at which only the players in S prefer x to y , that is,

$$\mathcal{C}^{\mathcal{P}}(S, x \succ y) := \{p \in \mathcal{P} \mid \mathcal{S}(p, x \succ y) = S\}.$$

Let us define $\mathcal{N}^{\mathcal{P}}(x \succ y)$, which indicates whether the players can express their opinions as $x \succ y$ or not, and similarly $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, as

$$\mathcal{N}^{\mathcal{P}}(x \succ y) := \begin{cases} \mathcal{N} & \text{if there exists a preference in } \Omega \text{ such that } x \succ y, \\ \emptyset & \text{otherwise,} \end{cases}$$

$$\mathcal{N}^{\mathcal{P}}(x \succ y \succ z) := \begin{cases} \mathcal{N} & \text{if there exists a preference in } \Omega \text{ such that } x \succ y \succ z, \\ \emptyset & \text{otherwise,} \end{cases}$$

for each distinct $x, y, z \in \mathcal{X}$ on a given $\mathcal{P} = \Omega^n$.

A *social welfare function* (on linear orderings) denoted by f , is a mapping that assigns an ordering on \mathcal{X} to a profile $p \in \mathcal{P}$, i.e., $f : \mathcal{P} \rightarrow L(\mathcal{X})$. We denote by $\succ^{f(p)}$ the social preference ordering on \mathcal{X} determined by f at profile $p \in \mathcal{P}$. A social welfare function is said to satisfy *unrestricted domain property* when $\Omega = L(\mathcal{X})$ holds. If there exists a player $i \in \mathcal{N}$, such that $x \succ_i^p y$ implies $x \succ^{f(p)} y$ for any pair of distinct alternatives $x, y \in \mathcal{X}$ and for any profile $p \in \mathcal{P}$, then the function is said to be *dictatorial* and i to be a *dictator*. A social welfare function satisfying the following two axioms is called an *Arrowian Social Welfare Function (ASWF)*.

Axiom 2.1 (Pareto Principle (PP)). If the property that

$$x \succ_i^p y \text{ for all } i \in \mathcal{N} \text{ implies } x \succ^{f(p)} y$$

holds for any pair of distinct alternatives $x, y \in \mathcal{X}$ and for any profile $p \in \mathcal{P}$, then the social welfare function f is said to have *Pareto principle*.

Axiom 2.2 (Independence of Irrelevant Alternatives (IIA)). If the property that

$$\succ_i^{p_1}|_{\{x,y\}} = \succ_i^{p_2}|_{\{x,y\}} \text{ for all } i \in \mathcal{N} \text{ implies } \succ^{f(p_1)}|_{\{x,y\}} = \succ^{f(p_2)}|_{\{x,y\}}$$

holds for any pair of distinct alternatives $x, y \in \mathcal{X}$ and for any pair of distinct profiles $p_1, p_2 \in \mathcal{P}$, then the social welfare function f is said to satisfy *independence of irrelevant alternatives*.

3. ALTERNATIVE PROOF OF INTEGER LINEAR INEQUALITY REPRESENTATION

This section shows an alternative way to prove the one-to-one correspondence theorem between an ASWF and an integer solution to the ASWF linear inequalities in Sethuraman et al. [11]. First of all, let ASWF f be given, and we introduce variable $d_S(x, y)$ for each pair of distinct alternatives $x, y \in \mathcal{X}$, and for each set S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$. The value of $d_S(x, y)$ is set by

$$(3.1) \quad d_S(x, y) := \begin{cases} 1 & \text{if } x \succ^{f(p)} y \text{ holds for all } p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y), \\ 0 & \text{otherwise,} \end{cases}$$

and we will give a certain integer linear inequality system to which the $d_S(x, y)$ constructed by this way becomes a solution. On the other hand, for a given solution to this inequality system, we construct f , a function which maps a profile into a binary relation on \mathcal{X} and is hopefully an ASWF, as follows: for each profile $p \in \mathcal{P}$ and for each distinct pair of alternatives $x, y \in \mathcal{X}$,

$$(3.2) \quad \begin{aligned} x \succ^{f(p)} y & \text{ if } d_{S(p, x \succ y)}(x, y) = 1, \text{ and} \\ x \not\succeq^{f(p)} y & \text{ otherwise.} \end{aligned}$$

We present below the inequality system with regard to $d_S(x, y)$.

Definition 3.1 (Integer Linear Inequality Representation of ASWFs).

(Integrality and IIA). For all pairs of distinct alternatives $x, y \in \mathcal{X}$, and for all sets S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$,

$$(3.3) \quad d_S(x, y) \in \{0, 1\}.$$

(PP). For all pairs of distinct alternatives $x, y \in \mathcal{X}$ such that $\mathcal{N}^{\mathcal{P}}(x \succ y) = \mathcal{N}$,

$$(3.4) \quad d_{\mathcal{N}}(x, y) = 1.$$

(Completeness and Antisymmetry). For all pairs of distinct alternatives $x, y \in \mathcal{X}$, and for all sets S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$,

$$(3.5) \quad d_S(x, y) + d_{\mathcal{N} \setminus S}(y, x) = 1.$$

(Transitivity). For all ordered triples (x, y, z) of distinct alternatives $x, y, z \in \mathcal{X}$, and for all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, C, U, V, W) is a partition of \mathcal{N} ,

$$(3.6) \quad d_{A \cup U \cup V}(x, y) + d_{B \cup U \cup W}(y, z) + d_{C \cup V \cup W}(z, x) \leq 2.$$

We note that there is a slight difference between the original ASWF formulation by Sethuraman et al. [11] and that by Definition 3.1. For any pair of distinct $x, y \in \mathcal{X}$, the set S takes between an empty set and the whole set \mathcal{N} of players in the former, while $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$ in the latter. The difference arises when $\mathcal{N}^{\mathcal{P}}(x \succ y) = \emptyset$ or $\mathcal{N}^{\mathcal{P}}(y \succ x) = \emptyset$, i.e., one alternative is always strictly preferred to the other for all preference orderings in the domain Ω . This situation is said that either (x, y) or (y, x) is a trivial pair (see Definition 4.4). In the original work a constant value is set to

the variables involving a trivial pair by convention, while we do not even enumerate such variables.

Theorem 3.2 (Sethuraman et al. [11, Theorem 1]). *Every solution to Integer Linear Inequality Representation of ASWFs corresponds to an ASWF on linear orderings and vice versa.*

We will present another proof of this theorem, which we believe will furnish us with a more insightful view. In the first stage of proving Theorem 3.2 we introduce another integer linear inequality system and show one-to-one correspondence between a solution to the system and an ASWF. We introduce the profile-dependent variable $d(p, x, y)$ and make an ASWF f and a vector d of $d(p, x, y)$'s correspond to each other by the following natural way: for each $p \in \mathcal{P}$ and for each distinct $x, y \in \mathcal{X}$,

$$(3.7) \quad \begin{aligned} x \succ^{f(p)} y &\text{ if and only if } d(p, x, y) = 1, \text{ and} \\ x \not\succeq^{f(p)} y &\text{ if and only if } d(p, x, y) = 0. \end{aligned}$$

We give the integer linear inequalities concerning the profile-dependent variable.

Definition 3.3 (Profile-dependent Integer Linear Inequality Representation of ASWFs).

(Integrality). For all profiles $p \in \mathcal{P}$, and for all pairs of distinct alternatives $x, y \in \mathcal{X}$,

$$(3.8) \quad d(p, x, y) \in \{0, 1\}.$$

(PP). For all profiles $p \in \mathcal{P}$, and for all pairs of distinct alternatives $x, y \in \mathcal{X}$, if $p \in \mathcal{C}^{\mathcal{P}}(\mathcal{N}, x \succ y)$ then

$$(3.9) \quad d(p, x, y) = 1.$$

(IIA). For all pairs of distinct profiles $p_1, p_2 \in \mathcal{P}$, and for all pairs of distinct alternatives $x, y \in \mathcal{X}$, if $p_1, p_2 \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$ for some S then

$$(3.10) \quad d(p_1, x, y) = d(p_2, x, y).$$

(Completeness and Antisymmetry). For all profiles $p \in \mathcal{P}$, and for all pairs of distinct alternatives $x, y \in \mathcal{X}$,

$$(3.11) \quad d(p, x, y) + d(p, y, x) = 1.$$

(Transitivity). For all profiles $p \in \mathcal{P}$, and for all triples of distinct alternatives $x, y, z \in \mathcal{X}$,

$$(3.12) \quad d(p, x, y) + d(p, y, z) \leq 1 + d(p, x, z).$$

Lemma 3.4. *Every solution to Profile-dependent Integer Linear Inequality Representation of ASWFs corresponds to an ASWF on linear orderings and vice versa.*

Proof. First we show that $d(p, x, y)$ constructed from a given ASWF f satisfies the inequalities in Definition 3.3. Needless to say (3.8) is satisfied. When $p \in \mathcal{C}^{\mathcal{P}}(\mathcal{N}, x \succ y)$, Axiom (PP) of an ASWF makes $x \succ^{f(p)} y$ hold, and therefore (3.9) is satisfied. By definition, the statement $p_1, p_2 \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$ implies $\mathcal{S}(p_1, x \succ y) = \mathcal{S}(p_2, x \succ y) = S$. Since the preferences of the players in p_1 and p_2 are all linear-ordered, both $\mathcal{S}(p_1, y \succ x)$ and $\mathcal{S}(p_2, y \succ x)$ are $\mathcal{N} \setminus S$. Hence $\succ_i^{p_1}|_{\{x, y\}} = \succ_i^{p_2}|_{\{x, y\}}$ holds for all $i \in \mathcal{N}$. Accordingly Axiom (IIA) guarantees $\succ^{f(p_1)}|_{\{x, y\}} = \succ^{f(p_2)}|_{\{x, y\}}$ and from that (3.10) is obtained. Completeness and antisymmetry of a social welfare function ensure (3.11), and by transitivity (3.12) is satisfied.

Next we let vector d be an arbitrary solution to (3.8)-(3.12) and let f be constructed by d . From (3.8) and (3.11), exactly one of either $x \succ^{f(p)} y$ or $y \succ^{f(p)} x$ holds for any profile $p \in \mathcal{P}$ and for any pair of distinct alternatives $x, y \in \mathcal{X}$, so that f is complete and antisymmetric. Suppose that f does not satisfy transitivity, i.e., $x \succ^{f(p)} y$, $y \succ^{f(p)} z$ and $x \not\succeq^{f(p)} z$ for some triple x, y and $z \in \mathcal{X}$. Then we have $d(p, x, y) = d(p, y, z) = 1$ and $d(p, x, z) = 0$, which violates (3.12). To show that f satisfies Axiom (PP), take an arbitrary pair of distinct alternatives $x, y \in \mathcal{X}$. Then for any $p \in \mathcal{P}$ such that $\mathcal{S}(p, x \succ y) = \mathcal{N}$, (3.9) and (3.11) indicate $d(p, x, y) = 1$ and $d(p, y, x) = 0$, which means $x \succ^{f(p)} y$. Choose an arbitrary pair of distinct $x, y \in \mathcal{X}$ as well as arbitrary distinct $p_1, p_2 \in \mathcal{P}$, and suppose that the statement $\succ_i^{p_1}|_{\{x, y\}} = \succ_i^{p_2}|_{\{x, y\}}$ holds for all $i \in \mathcal{N}$. It implies $p_1, p_2 \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$ for some S and $p_1, p_2 \in \mathcal{C}^{\mathcal{P}}(\mathcal{N} \setminus S, y \succ x)$. Then from (3.8) and (3.10), they hold that $x \succ^{f(p_1)} y$ if and only if $x \succ^{f(p_2)} y$ and $y \succ^{f(p_1)} x$ if and only if $y \succ^{f(p_2)} x$. Therefore we can say that f satisfies Axiom (IIA). Thus the lemma has been proved. \square

In the second stage for the proof of Theorem 3.2 we connect the profile-dependent representation with the original representation. Namely, let an arbitrary solution to (3.8)-(3.12) be given, and set $d_S(x, y)$ for each pair of distinct alternatives $x, y \in \mathcal{X}$, and for each set S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$ as

$$(3.13) \quad d_S(x, y) := \frac{\sum_{p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)} d(p, x, y)}{|\mathcal{C}^{\mathcal{P}}(S, x \succ y)|}.$$

In other words, we regard $d_S(x, y)$ as the ratio of x being socially preferred to y at the profiles such that all the players in S express their preferences as $x \succ y$ and the rest as $y \succ x$. We confirm that $\mathcal{C}^{\mathcal{P}}(S, x \succ y)$ is not empty for each pair of distinct alternatives $x, y \in \mathcal{X}$ and for each set S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$. When $\mathcal{N}^{\mathcal{P}}(x \succ y) = \emptyset$, S can only be \emptyset . In this case there does not exist a preference such that $x \succ y$. Therefore $\mathcal{S}(p, x \succ y) = \emptyset$ holds for all $p \in \mathcal{P}$, and equivalently $\mathcal{C}^{\mathcal{P}}(\emptyset, x \succ y) = \mathcal{P}$. When $\mathcal{N}^{\mathcal{P}}(x \succ y) = \mathcal{N}$ and $\mathcal{N}^{\mathcal{P}}(y \succ x) = \emptyset$, S can only be \mathcal{N} . This means that for all profiles every player strictly prefers x to y , thus $\mathcal{C}^{\mathcal{P}}(\mathcal{N}, x \succ y) \neq \emptyset$ holds. Each player can express $x \succ y$ or $y \succ x$ when both $\mathcal{N}^{\mathcal{P}}(x \succ y)$ and $\mathcal{N}^{\mathcal{P}}(y \succ x)$ are \mathcal{N} , so in this case $\mathcal{C}^{\mathcal{P}}(S, x \succ y)$ is nonempty for any $\emptyset \subseteq S \subseteq \mathcal{N}$. From this discussion we can say that $d_S(x, y)$ constructed by (3.13) is well-defined. Meanwhile, given a solution to Integer Linear Inequality Representation of ASWFs in the sense of Definition 3.1, we create $d(p, x, y)$ for each $p \in \mathcal{P}$ and for each distinct pair of alternatives $x, y \in \mathcal{X}$ as follows:

$$(3.14) \quad d(p, x, y) := d_{\mathcal{S}(p, x \succ y)}(x, y).$$

Lemma 3.5. *Every solution to Integer Linear Inequality Representation of ASWFs corresponds to a solution to Profile-dependent Integer Linear Inequality Representation of ASWFs and vice versa.*

Proof. Since $p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$ holds if and only if $\mathcal{S}(p, x \succ y) = S$ holds, $d(p, x, y)$ obtained from a given $d_S(x, y)$ by (3.14) returns the same $d_S(x, y)$ by (3.13). The converse also holds if (3.10) is assumed. Hence it suffices to show that a solution to (3.8)-(3.12) constructs a solution to (3.3)-(3.6) and a given $d_S(x, y)$ gives $d(p, x, y)$ which satisfies the constraints (3.8)-(3.12) in Definition 3.3.

Let an arbitrary solution to (3.8)-(3.12) be given. We take an arbitrary pair of distinct $x, y \in \mathcal{X}$, and an arbitrary but appropriate S . From (3.10) and (3.13), the statement

$$(3.15) \quad d_S(x, y) = d(p, x, y)$$




FIGURE 1. Correspondence among f , $d(p, x, y)$ and $d_S(x, y)$

holds for any $p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$. This equation along with (3.8) indicates (3.3). Consider the case where $S = \mathcal{N}^{\mathcal{P}}(x \succ y) = \mathcal{N}$, and we have (3.4) from (3.9). When a profile p is in $\mathcal{C}^{\mathcal{P}}(S, x \succ y)$ it is also a member of $\mathcal{C}^{\mathcal{P}}(\mathcal{N} \setminus S, y \succ x)$ because of the antisymmetry of a linear preference ordering. Therefore, (3.11) and (3.15) imply

$$d_S(x, y) + d_{\mathcal{N} \setminus S}(y, x) = d(p, x, y) + d(p, y, x) = 1,$$

which is (3.5). To show that (3.6) is satisfied, suppose not. Then for some distinct $x, y, z \in \mathcal{X}$ and for some partition (A, B, C, U, V, W) of \mathcal{N} with $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, and $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$,

$$d_{AUUV}(x, y) + d_{BUUW}(y, z) + d_{CUVUW}(z, x) > 2$$

holds. If player i is in A , there exists a preference ordering on which i can express his/her opinion as $x \succ z \succ y$. This is because $i \in A$ implies $\mathcal{N}^{\mathcal{P}}(x \succ z \succ y) = \mathcal{N}$, the existence of a preference with $x \succ z \succ y$. Likewise $i \in B$ means that i can choose a preference from Ω which is $y \succ x \succ z$, etc. Note also that each player is in exactly one of A, B, C, U, V , and W , and that we can choose a profile $p \in \mathcal{P}$ such that, for each $i \in \mathcal{N}$,

$$\begin{aligned} \text{if } i \in A \text{ then } x \succ_i^p z \succ_i^p y, & \quad \text{else if } i \in B \text{ then } y \succ_i^p x \succ_i^p z, \\ \text{else if } i \in C \text{ then } z \succ_i^p y \succ_i^p x, & \quad \text{else if } i \in U \text{ then } x \succ_i^p y \succ_i^p z, \\ \text{else if } i \in V \text{ then } z \succ_i^p x \succ_i^p y, & \quad \text{else if } i \in W \text{ then } y \succ_i^p z \succ_i^p x. \end{aligned}$$

For this profile, we have $p \in \mathcal{C}^{\mathcal{P}}(A \cup U \cup V, x \succ y)$, $p \in \mathcal{C}^{\mathcal{P}}(B \cup U \cup W, y \succ z)$, and $p \in \mathcal{C}^{\mathcal{P}}(C \cup V \cup W, z \succ x)$ hold. Then by (3.15), we have

$$d(p, x, y) + d(p, y, z) + d(p, z, x) > 2.$$

On the other hand by applying (3.11) to (3.12) we obtain

$$(3.16) \quad d(p, x, y) + d(p, y, z) + d(p, z, x) \leq 2,$$

consequently a contradiction.

Let a solution to Integer Linear Inequality Representation of ASWFs in the sense of Definition 3.1 be given, and the value of $d(p, x, y)$ be set by (3.14). We show that this $d(p, x, y)$ satisfies (3.8)-(3.12). The constructed $d(p, x, y)$ by this way satisfies (3.10) since the statement $p_1, p_2 \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$ is equivalent to $\mathcal{S}(p_1, x \succ y) = \mathcal{S}(p_2, x \succ y) = S$.

In the case where $S = \mathcal{N}$, (3.9) is implied by (3.4). The set $\mathcal{S}(p, x \succ y)$ lies between $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x)$ and $\mathcal{N}^{\mathcal{P}}(x \succ y)$, hence we have (3.8) from (3.3). Antisymmetry of preference orderings in the common preference domain framework guarantees $\mathcal{S}(p, y \succ x) = \mathcal{N} \setminus \mathcal{S}(p, x \succ y)$, and this together with (3.5) assures us

$$d(p, x, y) + d(p, y, x) = d_{\mathcal{S}(p, x \succ y)}(x, y) + d_{\mathcal{S}(p, y \succ x)}(y, x) = 1,$$

which is (3.11). Let us suppose that (3.12) does not hold for some $p \in \mathcal{P}$ and for some distinct alternatives $x, y, z \in \mathcal{X}$. Then (3.16) is also violated by these p, x, y , and z , i.e.,

$$d(p, x, y) = 1, d(p, y, z) = 1, \text{ and } d(p, z, x) = 1.$$

It means, by (3.14), that

$$d_{\mathcal{S}(p, x \succ y)}(x, y) = 1, d_{\mathcal{S}(p, y \succ z)}(y, z) = 1, \text{ and } d_{\mathcal{S}(p, z \succ x)}(z, x) = 1,$$

respectively. Regarding the triple $\{x, y, z\}$, there are six patterns of preference orderings at this p . We partition the whole set \mathcal{N} of players into the six types as follows:

$$\begin{aligned} \bar{A} &:= \{i \in N \mid x \succ_i^p z \succ_i^p y\}, & \bar{B} &:= \{i \in N \mid y \succ_i^p x \succ_i^p z\}, \\ \bar{C} &:= \{i \in N \mid z \succ_i^p y \succ_i^p x\}, & \bar{U} &:= \{i \in N \mid x \succ_i^p y \succ_i^p z\}, \\ \bar{V} &:= \{i \in N \mid z \succ_i^p x \succ_i^p y\}, & \bar{W} &:= \{i \in N \mid y \succ_i^p z \succ_i^p x\}. \end{aligned}$$

Since $\mathcal{S}(p, x \succ y) = \bar{A} \cup \bar{U} \cup \bar{V}$, $\mathcal{S}(p, y \succ z) = \bar{B} \cup \bar{U} \cup \bar{W}$, and $\mathcal{S}(p, z \succ x) = \bar{C} \cup \bar{V} \cup \bar{W}$ hold, we have

$$d_{\bar{A} \cup \bar{U} \cup \bar{V}}(x, y) + d_{\bar{B} \cup \bar{U} \cup \bar{W}}(y, z) + d_{\bar{C} \cup \bar{V} \cup \bar{W}}(z, x) = 3.$$

This contradicts (3.6) because the sets $\bar{A}, \bar{B}, \bar{C}, \bar{U}, \bar{V}$, and \bar{W} are all disjoint, their union coincides with \mathcal{N} , and \bar{A} is a subset of $\mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\bar{B} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, etc. \square

Combining Lemma 3.4 and Lemma 3.5 we have the following proof.

Proof of Theorem 3.2. By Lemma 3.4 and Lemma 3.5 along with the transformation procedures (3.7), (3.13), and (3.14), we have the one-to-one correspondence between an ASWF and a solution to Integer Linear Inequality Representation of ASWFs. \square

We note that the transitivity inequalities (3.6) ‘for all “ordered” triples’ can be substituted for those ‘for all “unordered” triples,’ which contributes to a reduction in the number of constraints.

Definition 3.6 (Transitivity on Unordered Triples). For all unordered triples $\{x, y, z\}$ of distinct alternatives $x, y, z \in \mathcal{X}$, and for all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, C, U, V, W) is a partition of \mathcal{N} ,

$$(3.17) \quad 1 \leq d_{AUUV}(x, y) + d_{BUUW}(y, z) + d_{CUVW}(z, x) \leq 2.$$

Theorem 3.7. When (3.5) is assumed, (3.6) is equivalent to (3.17).

Proof. Let $\{x, y, z\}$ be an arbitrary unordered triple of distinct alternatives $x, y, z \in \mathcal{X}$. We evaluate (3.6) for all permutations of $\{x, y, z\}$, i.e., (x, y, z) , (y, z, x) , (z, x, y) , (x, z, y) , (z, y, x) , and (y, x, z) .

Case 1: (x, y, z) . For all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, C, U, V, W) is a partition of \mathcal{N} ,

$$(3.18) \quad d_{AUUV}(x, y) + d_{BUUW}(y, z) + d_{CUVW}(z, x) \leq 2.$$

Case 2: (y, z, x) . For all sets $\bar{A} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\bar{B} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\bar{C} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\bar{U} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, $\bar{V} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $\bar{W} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, such that $(\bar{A}, \bar{B}, \bar{C}, \bar{U}, \bar{V}, \bar{W})$ is a partition of \mathcal{N} , it clearly holds that

$$(3.19) \quad d_{\bar{A}\bar{U}\bar{U}\bar{V}}(y, z) + d_{\bar{B}\bar{U}\bar{U}\bar{W}}(z, x) + d_{\bar{C}\bar{U}\bar{V}\bar{U}\bar{W}}(x, y) \leq 2.$$

By letting $\bar{A} = B$, $\bar{B} = C$, $\bar{C} = A$, $\bar{U} = W$, $\bar{V} = U$, and $\bar{W} = V$, we see that (3.19) is the same as (3.18).

Case 3: (z, x, y) . For all sets $\bar{A} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\bar{B} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\bar{C} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\bar{U} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $\bar{V} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, $\bar{W} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, such that $(\bar{A}, \bar{B}, \bar{C}, \bar{U}, \bar{V}, \bar{W})$ is a partition of \mathcal{N} ,

$$(3.20) \quad d_{\bar{A}\bar{U}\bar{U}\bar{V}}(z, x) + d_{\bar{B}\bar{U}\bar{U}\bar{W}}(x, y) + d_{\bar{C}\bar{U}\bar{V}\bar{U}\bar{W}}(y, z) \leq 2.$$

Setting $\bar{A} = C$, $\bar{B} = A$, $\bar{C} = B$, $\bar{U} = V$, $\bar{V} = W$, and $\bar{W} = U$ tells us that (3.20) is also equivalent to (3.18).

Case 4: (x, z, y) . For all sets $\hat{A} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $\hat{B} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $\hat{C} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, $\hat{U} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\hat{V} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\hat{W} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, such that $(\hat{A}, \hat{B}, \hat{C}, \hat{U}, \hat{V}, \hat{W})$ is a partition of \mathcal{N} ,

$$(3.21) \quad d_{\hat{A}\hat{U}\hat{U}\hat{V}}(x, z) + d_{\hat{B}\hat{U}\hat{U}\hat{W}}(z, y) + d_{\hat{C}\hat{U}\hat{V}\hat{U}\hat{W}}(y, x) \leq 2.$$

We let $\hat{A} = U$, $\hat{B} = V$, $\hat{C} = W$, $\hat{U} = A$, $\hat{V} = B$, $\hat{W} = C$, then (3.21) is as this: for all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, C, U, V, W) is a partition of \mathcal{N} ,

$$(3.22) \quad d_{U\cup A\cup B}(x, z) + d_{V\cup A\cup C}(z, y) + d_{W\cup B\cup C}(y, x) \leq 2.$$

By applying (3.5) to all the variables in (3.22) we obtain the following inequality: for all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, C, U, V, W) is a partition of \mathcal{N} ,

$$d_{A\cup U\cup V}(x, y) + d_{B\cup U\cup W}(y, z) + d_{C\cup V\cup W}(z, x) \geq 1.$$

Case 5: (z, y, x) . For all sets $\bar{A} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $\bar{B} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, $\bar{C} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $\bar{U} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\bar{V} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\bar{W} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, such that $(\bar{A}, \bar{B}, \bar{C}, \bar{U}, \bar{V}, \bar{W})$ is a partition of \mathcal{N} ,

$$(3.23) \quad d_{\bar{A}\bar{U}\bar{U}\bar{V}}(z, y) + d_{\bar{B}\bar{U}\bar{U}\bar{W}}(y, x) + d_{\bar{C}\bar{U}\bar{V}\bar{U}\bar{W}}(x, z) \leq 2.$$

We see that (3.23) is equivalent to (3.21) by setting $\bar{A} = \hat{B}$, $\bar{B} = \hat{C}$, $\bar{C} = \hat{A}$, $\bar{U} = \hat{W}$, $\bar{V} = \hat{U}$, and $\bar{W} = \hat{V}$.

Case 6: (y, x, z) . For all sets $\bar{A} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, $\bar{B} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $\bar{C} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $\bar{U} \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\bar{V} \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\bar{W} \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, such that $(\bar{A}, \bar{B}, \bar{C}, \bar{U}, \bar{V}, \bar{W})$ is a partition of \mathcal{N} ,

$$(3.24) \quad d_{\bar{A}\bar{U}\bar{U}\bar{V}}(y, x) + d_{\bar{B}\bar{U}\bar{U}\bar{W}}(x, z) + d_{\bar{C}\bar{U}\bar{V}\bar{U}\bar{W}}(z, y) \leq 2.$$

Again we have (3.21) if we let $\bar{A} = \hat{C}$, $\bar{B} = \hat{A}$, $\bar{C} = \hat{B}$, $\bar{U} = \hat{V}$, $\bar{V} = \hat{W}$, and $\bar{W} = \hat{U}$ in (3.24). \square

Corollary 3.8. *Every solution to (3.3)-(3.5) and (3.17) corresponds to an ASWF on linear orderings and vice versa.*

4. POLYHEDRAL STRUCTURE OF LINEAR INEQUALITIES

This section studies some polyhedral structure determined by the linear inequalities (3.4)-(3.6) along with the nonnegativity of the variables in Integer Linear Inequality Representation of ASWFs. We introduce the weakly nonisolated condition on the preference domain and present that all the extreme points of the polytope are integers when the domain is single-peaked, single-caved, or the domain on which each triple of alternatives contains an alternative that cannot be medium, and it satisfies the weakly nonisolated condition.

Definition 4.1 (Single-peakedness). Let \triangleright be a linear ordering on \mathcal{X} , i.e., $\triangleright \in L(\mathcal{X})$. The domain Ω is called *single-peaked with respect to* \triangleright if Ω is a subset of $\Omega^P(\triangleright)$ where

$$\Omega^P(\triangleright) := \{ \succ \in L(\mathcal{X}) \mid \text{for every } x, y, z \in \mathcal{X} \text{ with } x \triangleright y \triangleright z, y \succ x \text{ or } y \succ z \text{ holds} \}.$$

We refer to the linear ordering \triangleright as the *reference linear ordering*.

The class of single-peaked domains is well known and seen in Arrow [2], Sen [10], and other many books and papers. Concerning the polyhedral structure on single-peaked domains the following is claimed in Sethuraman et al. [11].

Claim 4.2 (Sethuraman et al. [11, Theorem 10]). *When Ω is single-peaked (with respect to $\triangleright \in L(\mathcal{X})$) the set of nonnegative solutions satisfying (3.4)-(3.6) is an integral polytope, i.e., all the extreme points are integer vectors.*

We discuss here an example of the ASWF linear inequality formulation and its polytope on a single-peaked domain.

Example 4.3. Let $n = 2$, $\mathcal{X} = \{u, v, w, x, y, z\}$, $\Omega = \{u \succ v \succ w \succ x \succ y \succ z, z \succ y \succ x \succ w \succ v \succ u\}$, and the reference linear ordering \triangleright be $u \triangleright v \triangleright w \triangleright x \triangleright y \triangleright z$. The domain Ω is obviously single-peaked with respect to \triangleright . We show below its ASWF linear inequalities (3.4), (3.5), and (3.17) that determine a polytope. Note that we choose here an unordered triple $\{a, b, c\}$ with $a \triangleright b \triangleright c$ to enumerate all the inequalities of (3.17) for arbitrary distinct alternatives $a, b, c \in \mathcal{X}$. The sets A, B, V , and W in (3.17) must always be empty since only the preferences with $a \succ b \succ c$ and $c \succ b \succ a$ are admissible.

For all pairs of distinct alternatives $a, b \in \mathcal{X}$,

$$d_{\mathcal{N}}(a, b) = 1, d_{\emptyset}(a, b) = 0, d_{\{2\}}(a, b) = 1 - d_{\{1\}}(b, a).$$

(4.1) For all unordered triple $\{a, b, c\}$ of alternatives $a, b, c \in \mathcal{X}$ such that $a \triangleright b \triangleright c$, and for all disjoint sets U, C such that $C \cup U = \mathcal{N}$,

$$1 \leq d_U(a, b) + d_U(b, c) + d_C(c, a) \leq 2.$$

To this linear inequality system, a fractional solution

(4.2)

$$\begin{aligned} d_{\{1\}}(u, v) &= 0.5, & d_{\{1\}}(u, w) &= 0, & d_{\{1\}}(u, x) &= 0.5, & d_{\{1\}}(u, y) &= 0.5, & d_{\{1\}}(u, z) &= 0, \\ d_{\{1\}}(v, u) &= 0, & d_{\{1\}}(v, w) &= 0, & d_{\{1\}}(v, x) &= 0.5, & d_{\{1\}}(v, y) &= 0, & d_{\{1\}}(v, z) &= 0.5, \\ d_{\{1\}}(w, u) &= 0, & d_{\{1\}}(w, v) &= 0, & d_{\{1\}}(w, x) &= 0.5, & d_{\{1\}}(w, y) &= 0.5, & d_{\{1\}}(w, z) &= 0.5, \\ d_{\{1\}}(x, u) &= 0, & d_{\{1\}}(x, v) &= 0, & d_{\{1\}}(x, w) &= 0, & d_{\{1\}}(x, y) &= 0, & d_{\{1\}}(x, z) &= 0, \\ d_{\{1\}}(y, u) &= 0, & d_{\{1\}}(y, v) &= 0, & d_{\{1\}}(y, w) &= 0, & d_{\{1\}}(y, x) &= 0, & d_{\{1\}}(y, z) &= 0.5, \\ d_{\{1\}}(z, u) &= 0, & d_{\{1\}}(z, v) &= 0, & d_{\{1\}}(z, w) &= 0, & d_{\{1\}}(z, x) &= 0, & d_{\{1\}}(z, y) &= 0, \\ \text{and } d_{\mathcal{N}}(a, b) &= 1, & d_{\emptyset}(a, b) &= 0, & d_{\{2\}}(a, b) &= 1 - d_{\{1\}}(b, a) \end{aligned}$$

for all pairs of distinct alternatives $a, b \in \mathcal{X}$,

can be obtained and is seen to be the unique solution to the following equalities:

$$\begin{aligned}
& d_{\{1\}}(u, w) = 0, \quad d_{\{1\}}(u, z) = 0, \quad d_{\{1\}}(v, u) = 0, \quad d_{\{1\}}(v, w) = 0, \quad d_{\{1\}}(v, y) = 0, \\
& d_{\{1\}}(w, u) = 0, \quad d_{\{1\}}(w, v) = 0, \quad d_{\{1\}}(x, u) = 0, \quad d_{\{1\}}(x, v) = 0, \quad d_{\{1\}}(x, w) = 0, \\
& d_{\{1\}}(x, y) = 0, \quad d_{\{1\}}(x, z) = 0, \quad d_{\{1\}}(y, u) = 0, \quad d_{\{1\}}(y, v) = 0, \quad d_{\{1\}}(y, w) = 0, \\
& d_{\{1\}}(y, x) = 0, \quad d_{\{1\}}(z, u) = 0, \quad d_{\{1\}}(z, v) = 0, \quad d_{\{1\}}(z, w) = 0, \quad d_{\{1\}}(z, x) = 0, \\
& d_{\{1\}}(z, y) = 0, \\
& d_{\{1\}}(u, v) + d_{\{1\}}(v, y) + d_{\{2\}}(y, u) = 1, \quad d_{\{1\}}(u, w) + d_{\{1\}}(w, x) + d_{\{2\}}(x, u) = 1, \\
& d_{\{1\}}(u, w) + d_{\{1\}}(w, y) + d_{\{2\}}(y, u) = 1, \quad d_{\{1\}}(u, x) + d_{\{1\}}(x, y) + d_{\{2\}}(y, u) = 1, \\
& d_{\{1\}}(v, w) + d_{\{1\}}(w, x) + d_{\{2\}}(x, v) = 1, \quad d_{\{1\}}(v, w) + d_{\{1\}}(w, z) + d_{\{2\}}(z, y) = 1, \\
& d_{\{1\}}(v, x) + d_{\{1\}}(x, z) + d_{\{2\}}(z, v) = 1, \\
& d_{\{1\}}(u, v) + d_{\{1\}}(v, z) + d_{\{2\}}(z, u) = 2, \quad d_{\{1\}}(u, y) + d_{\{1\}}(y, z) + d_{\{2\}}(z, u) = 2, \\
& \text{and } d_{\mathcal{N}}(a, b) = 1, \quad d_{\emptyset}(a, b) = 0, \quad d_{\{2\}}(a, b) = 1 - d_{\{1\}}(b, a)
\end{aligned}$$

for all pairs of distinct alternatives $a, b \in \mathcal{X}$.

Thus the fractional solution given by (4.2) is an extreme point of the set determined by the linear inequality system (4.1). This is a counter-example to Claim 4.2.

We impose an additional condition on a single-peaked domain and make the polytope induced by the inequalities on the domain integral. For the introduction of the new condition let us present the commonly known concept of a trivial pair of alternatives on admissible preferences, and then we define triviality on an unordered triple of distinct alternatives for convenience.

Definition 4.4 (Trivial Pair). An ordered pair (x, y) of distinct alternatives $x, y \in \mathcal{X}$ is said to be a *trivial pair on Ω* if $x \succ y$ holds for all preference orderings in Ω .

Definition 4.5 (Triviality over an Unordered Triple). For an unordered triple of distinct alternatives $x, y, z \in \mathcal{X}$, we call that $\{x, y, z\}$ has a *trivial pair on Ω* if at least one of (x, y) , (y, x) , (x, z) , (z, x) , (y, z) , and (z, y) is a trivial pair.

Then we refer to the notion of an isolated triple, which is originally proposed by Sethuraman et al. [11], and define nonisolation on an unordered triple.

Definition 4.6 (Isolated Triple). An ordered triple (x, y, z) of distinct alternatives $x, y, z \in \mathcal{X}$, is said to be an *isolated triple on Ω* if there exists a preference ordering in Ω with $x \succ y \succ z$ and there does not exist with $y \succ z \succ x$ or $z \succ x \succ y$.

Definition 4.7 (Nonisolation over an Unordered Triple). For an unordered triple of distinct alternatives $x, y, z \in \mathcal{X}$, we say that $\{x, y, z\}$ has *nonisolation on Ω* if none of (x, y, z) , (y, z, x) , (z, x, y) , (x, z, y) , (z, y, x) , and (y, x, z) is an isolated triple.

From the above definitions we finally introduce the condition on the preference domain that we call weak nonisolation and show the corrected version of Claim 4.2.

Definition 4.8 (Weak Nonisolation). The domain Ω is said to be *weakly nonisolated* if every unordered triple in \mathcal{X} has either a trivial pair or nonisolation on Ω .

Theorem 4.9. *When Ω is single-peaked with respect to the reference linear ordering $\triangleright \in L(\mathcal{X})$ and weakly nonisolated, the set of nonnegative solutions satisfying (3.4)-(3.6) is an integral polytope.*

Proof. We follow the technique of proof in Sethuraman et al. [11]: given an arbitrary nonnegative solution to the linear inequalities, we round it to an integer by a certain procedure, show that the rounded solution still satisfies the inequalities, and see that these facts guarantee the integrality of the polytope. Let \mathcal{O}^P be the polytope we are to

discuss, that is, the set of nonnegative solutions satisfying (3.4), (3.5), and (3.17) instead of (3.6) on the given $\mathcal{P} = \Omega^n$. Let d be an arbitrary vector of $d_S(x, y)$'s with $d \in \mathcal{O}^{\mathcal{P}}$. We generate a random number Z from the uniform distribution between 0 and 1. We round the vector d to a 0-1 vector, say d' , by the following way: for each pair of distinct alternatives $a, b \in \mathcal{X}$, and for each S with $\mathcal{N}^{\mathcal{P}}(a \succ b) \setminus \mathcal{N}^{\mathcal{P}}(b \succ a) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(a \succ b)$,

$$\text{if } a \triangleright b, d'_S(a, b) := \begin{cases} 1 & \text{if } d_S(a, b) > Z, \\ 0 & \text{otherwise,} \end{cases} \quad \text{else, } d'_S(a, b) := \begin{cases} 1 & \text{if } d_S(a, b) \geq 1 - Z, \\ 0 & \text{otherwise.} \end{cases}$$

We see that d' trivially satisfies (3.4) for any Z . Because $d_S(x, y) + d_{\mathcal{N} \setminus S}(y, x) = 1$ for an arbitrary pair of distinct alternatives $x, y \in \mathcal{X}$ and for an arbitrary appropriate S , $d_S(x, y) > Z$ holds if and only if $d_{\mathcal{N} \setminus S}(y, x) < 1 - Z$ holds and $d_S(x, y) \leq Z$ if and only if $d_{\mathcal{N} \setminus S}(y, x) \geq 1 - Z$. Exactly either one of $d_S(x, y)$ or $d_{\mathcal{N} \setminus S}(y, x)$ is 1 and the other is 0 when rounded, which means that (3.5) is also satisfied for d' .

Take an arbitrary triple of distinct alternatives $x, y, z \in \mathcal{X}$ with $x \triangleright y \triangleright z$, and consider (3.17) for $\{x, y, z\}$. By the single-peakedness of the domain with respect to \triangleright , there does not exist a preference in Ω such that $x \succ z \succ y$ or $z \succ x \succ y$. The sets $\mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$ and $\mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$ are empty, and (3.17) for $\{x, y, z\}$ can be written without the sets A and V : for all sets $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $U \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (B, C, U, W) is a partition of \mathcal{N} ,

$$(4.3) \quad 1 \leq d_U(x, y) + d_{B \cup U \cup W}(y, z) + d_{C \cup W}(z, x) \leq 2.$$

Case 1: $\{x, y, z\}$ has a trivial pair. When (x, y) is trivial, there does not exist a preference with $y \succ x$, which indicates that $\mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are all empty. Since (B, C, U, W) is a partition of \mathcal{N} , U must be \mathcal{N} , and we have $d_U(x, y) = 1$, $d_{B \cup U \cup W}(y, z) = 1$, and $d_{C \cup W}(z, x) = d_{\emptyset}(z, x) = 0$ from (3.4) and (3.5). Then the rounded solution satisfies (4.3), for 1 is rounded to 1 and 0 to 0 for any Z .

When (y, x) is trivial, $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z) = \emptyset$, and $d_U(x, y) = d_{\emptyset}(x, y) = 0$. Then the rounded solution d' trivially satisfies the ‘‘equal to or less than 2’’ part of (4.3). The rest is to show that

$$1 \leq d_{B \cup W}(y, z) + d_{C \cup W}(z, x)$$

is satisfied even after d is rounded. We see that if $d_{B \cup W}(y, z)$ is equal to or less than Z then the other variable is equal to or more than $1 - Z$, and that if $d_{C \cup W}(z, x)$ is less than $1 - Z$ the rest exceeds Z . Therefore d' satisfies this inequality from the rounding rule as well as $y \triangleright z$ and $x \triangleright z$.

When (x, z) is trivial, $\mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$ and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are the empty set. Accordingly $B \cup U = \mathcal{N}$ holds. This indicates $d_{B \cup U \cup W}(y, z) = 1$ and $d_{C \cup W}(z, x) = 0$, so the rounded d' clearly satisfies (4.3).

When (z, x) is trivial, it means $\mathcal{N}^{\mathcal{P}}(y \succ x \succ z) = \mathcal{N}^{\mathcal{P}}(x \succ y \succ z) = \emptyset$. Then $d_U(x, y) = 0$, and $d_{C \cup W}(z, x) = 1$ due to $C \cup W = \mathcal{N}$. In this case d' satisfies (4.3) again.

When (y, z) is trivial, $C \subseteq \mathcal{N}^{\mathcal{P}}(z \succ y \succ x) = \emptyset$, and $d_{B \cup U \cup W}(y, z) = 1$ since $B \cup U \cup W$ must be \mathcal{N} . The rounded solution trivially satisfies the ‘‘equal to or more than 1’’ part of (4.3). The rest is to prove that

$$d_U(x, y) + d_W(z, x) \leq 1$$

is kept satisfied after the rounding procedure. Assume here that $d_U(x, y)$ is more than Z , then the other is less than $1 - Z$. When $d_W(z, x) \geq 1 - Z$ holds, $d_U(x, y)$ is equal or less than Z . Hence by $x \triangleright y$ and $x \triangleright z$ this inequality is satisfied for d' .

When (z, y) is trivial, $\mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are empty. This means $B = U = W = \emptyset$ and $C = \mathcal{N}$. Accordingly we obtain $d_U(x, y) = d_{B \cup U \cup W}(y, z) = 0$ and $d_{C \cup W}(z, x) = 1$. The variables do not change if rounded.

Case 2: $\{x, y, z\}$ does not have a trivial pair. From the single-peakedness of Ω there is not a preference with $z \succ x \succ y$ or $x \succ z \succ y$. Then there exist preferences in Ω with $x \succ y \succ z$, $z \succ y \succ x$, respectively, otherwise (y, x) or (y, z) is trivial. Since $\{x, y, z\}$ is nonisolated, a preference with $y \succ z \succ x$ exists in Ω , and also a preference such that $y \succ x \succ z$. Therefore, $\mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are all equal to \mathcal{N} , and (4.3) holds for any partition (B, C, U, W) of \mathcal{N} . Take an arbitrary partition (B, C, U, W) , and let $W' = C \cup W$, $C' = \emptyset$. Because (B, C', U, W') is also a partition of \mathcal{N} , d satisfies the following:

$$\begin{aligned} 2 &\geq d_U(x, y) + d_{B \cup U \cup W'}(y, z) + d_{C' \cup W'}(z, x) = d_U(x, y) + d_{B \cup U \cup C \cup W}(y, z) + d_{C \cup W}(z, x) \\ &= d_U(x, y) + 1 + d_{C \cup W}(z, x). \end{aligned}$$

By rounding $d_U(x, y)$ and $d_{C \cup W}(z, x)$ in this inequality, we have

$$d'_U(x, y) + d'_{C \cup W}(z, x) \leq 1$$

from the rounding procedure together with $x \triangleright y$ and $x \triangleright z$. We see that d' satisfies the “equal to or less than 2” part of (4.3) whether $d_{B \cup U \cup W}(y, z)$ is rounded to 1 or 0. Regarding the “equal to or more than 1” part of (4.3), let $B' = B \cup U$, $U' = \emptyset$, and (B', C, U', W) is also a partition of \mathcal{N} . Hence,

$$1 \leq d_{U'}(x, y) + d_{B' \cup U' \cup W}(y, z) + d_{C \cup W}(z, x) = d_{B \cup U \cup W}(y, z) + d_{C \cup W}(z, x)$$

holds. We round the vector d in this inequality, then

$$d'_{B \cup U \cup W}(y, z) + d'_{C \cup W}(z, x) \geq 1$$

is obtained. This observation implies that d' satisfies the “equal to or more than 1” part of (4.3) regardless of the rounded value of $d_U(x, y)$. Hence we have confirmed that for any $d \in \mathcal{O}^{\mathcal{P}}$ the rounded vector d' is also in $\mathcal{O}^{\mathcal{P}}$.

Now we are ready to consider our main problem. Suppose that the polytope $\mathcal{O}^{\mathcal{P}}$, the set of nonnegative solutions which satisfies (3.4), (3.5), and (3.17), has an extreme point which is not an integer. Here we let \bar{d} be such a point. Since \bar{d} is an extreme point of $\mathcal{O}^{\mathcal{P}}$ there exists a cost vector c such that \bar{d} becomes the unique optimal solution to the linear programming problem:

$$(P) \quad \begin{cases} \min & c^\top d \\ \text{s.t.} & d \in \mathcal{O}^{\mathcal{P}} \end{cases}$$

where c^\top denotes the transpose of c . We generate a random number Z and round \bar{d} by the procedure proposed above. The rounded vector, denoted by $\bar{d}^{Z'}$, is still in $\mathcal{O}^{\mathcal{P}}$ as we have discussed. Now \bar{d} is assumed to be the unique optimal solution to (P), then

$$(4.4) \quad c^\top \bar{d}^{Z'} > c^\top \bar{d}$$

holds for any Z . Meanwhile, the expected value of each component of $\bar{d}^{Z'}$ is

$$\begin{aligned} \text{if } a \triangleright b, \quad & E(\bar{d}^{Z'}_S(a, b)) = 0 \times P(Z \geq \bar{d}_S(a, b)) + 1 \times P(Z < \bar{d}_S(a, b)) = \bar{d}_S(a, b), \\ \text{else,} \quad & E(\bar{d}^{Z'}_S(a, b)) = 0 \times P(Z < 1 - \bar{d}_S(a, b)) + 1 \times P(Z \geq 1 - \bar{d}_S(a, b)) = \bar{d}_S(a, b), \end{aligned}$$

for each $a, b \in \mathcal{X}$ and for each S which lies between $\mathcal{N}^{\mathcal{P}}(a \succ b) \setminus \mathcal{N}^{\mathcal{P}}(b \succ a)$ and $\mathcal{N}^{\mathcal{P}}(a \succ b)$, since Z is uniformly distributed between 0 and 1. This fact implies

$$E(c^\top \bar{d}^Z) = c^\top \bar{d},$$

contradicting (4.4). Thus all the extreme points of $\mathcal{O}^{\mathcal{P}}$ are integers. \square

We give another type of preference domains, called single-caved domains, and show that such domains along with weakly nonisolated condition also form integral polytopes.

Definition 4.10 (Single-cavedness). The domain Ω is called *single-caved with respect to* $\triangleright \in L(\mathcal{X})$ if Ω is a subset of $\Omega^C(\triangleright)$ where

$$\Omega^C(\triangleright) := \{ \succ \in L(\mathcal{X}) \mid \text{for every } x, y, z \in \mathcal{X} \text{ with } x \triangleright y \triangleright z, x \succ y \text{ or } z \succ y \text{ holds} \}.$$

Theorem 4.11. *When Ω is single-caved with respect to $\triangleright \in L(\mathcal{X})$ and weakly nonisolated, the set of nonnegative solutions satisfying (3.4)-(3.6) is an integral polytope.*

Proof. We can apply the same proof technique as the case where Ω is single-peaked to this case, so we omit the proof. \square

Single-peakedness is interpreted that one of three alternatives cannot be worst among the three while single-cavedness is that one of three cannot be best. We define the situation where one alternative cannot be medium among three alternatives. Then we derive integral polytopes on the domains that have such property.

Definition 4.12 (Cannot-be-medium Property). For a triple of distinct alternatives $x, y, z \in \mathcal{X}$, it is said that *y cannot be medium among (x, y, z) on Ω* if there does not exist a preference ordering in Ω such that $x \succ y \succ z$ or $z \succ y \succ x$.

Theorem 4.13. *When each triple of alternatives contains an alternative that cannot be medium on Ω and Ω is weakly nonisolated, the set of nonnegative solutions satisfying (3.4)-(3.6) is an integral polytope.*

Proof. Again we apply the same technique as when Ω is single-peaked to this proof, except that the rounding rule is changed as follows: for each pair of distinct alternatives $a, b \in \mathcal{X}$, and for each S with $\mathcal{N}^{\mathcal{P}}(a \succ b) \setminus \mathcal{N}^{\mathcal{P}}(b \succ a) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(a \succ b)$,

$$\text{if } \{1\} \in S, d'_S(a, b) := \begin{cases} 1 & \text{if } d_S(a, b) > Z, \\ 0 & \text{otherwise,} \end{cases} \quad \text{else, } d'_S(a, b) := \begin{cases} 1 & \text{if } d_S(a, b) \geq 1 - Z, \\ 0 & \text{otherwise.} \end{cases}$$

Given an arbitrary vector $d \in \mathcal{O}^{\mathcal{P}}$, the rounded d' clearly satisfies (3.4) for any Z , and also (3.5) since player 1 is always a member of either S or $\mathcal{N} \setminus S$ but not both. Now it suffices to show that d' satisfies (3.17). Let us take an arbitrary triple of distinct alternatives $x, y, z \in \mathcal{X}$, and consider (3.17) for $\{x, y, z\}$ with y cannot be medium. Then $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z) = \mathcal{N}^{\mathcal{P}}(z \succ y \succ x) = \emptyset$ holds, and (3.17) for $\{x, y, z\}$ reduces to this: for all sets $A \subseteq \mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $V \subseteq \mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$, such that (A, B, V, W) is a partition of \mathcal{N} ,

$$(4.5) \quad 1 \leq d_{A \cup V}(x, y) + d_{B \cup W}(y, z) + d_{V \cup W}(z, x) \leq 2.$$

Case 1: $\{x, y, z\}$ has a trivial pair. When (x, y) is trivial, B and W must be empty and $A \cup V = \mathcal{N}$ holds. Then $d_{A \cup V}(x, y) = d_{\mathcal{N}}(x, y) = 1$ as well as $d_{B \cup W}(y, z) = d_{\emptyset}(y, z) = 0$ holds. Recall that 1 is rounded to 1 and 0 to 0 for any Z , and we see that the rounded d' satisfies (4.5) regardless of the value of $d'_{V \cup W}(z, x)$.

When (y, x) is trivial, $A = V = \emptyset$ and the union of B and W is the whole set of players. This time $d_{A \cup V}(x, y) = 0$ and $d_{B \cup W}(y, z) = 1$ hold. Again (4.5) is still satisfied after d is rounded.

When (x, z) is trivial, both $\mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$ and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are empty, and accordingly V and W are. Then $d_{V \cup W}(z, x) = 0$ and the rounded d' trivially satisfies the “equal to or less than 2” part of (4.5). We round the rest inequality:

$$1 \leq d_A(x, y) + d_B(y, z).$$

Player 1 is in exactly either A or B , so we see that d' still satisfies this inequality.

When (z, x) is trivial, it means $A = B = \emptyset$ and $d_{V \cup W}(z, x) = d_{\mathcal{N}}(z, x) = 1$. Hence the “equal to or more than 1” part of (4.5) is cleared, and

$$d_V(x, y) + d_W(y, z) \leq 1$$

is still satisfied after the rounding, for player 1 is a member of either V or W but not both.

When (y, z) is trivial, $\mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$ and $\mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$ are empty. The set of players \mathcal{N} is partitioned into B and W , which is the same case as (y, x) -trivial.

When (z, y) is trivial, $B \subseteq \mathcal{N}^{\mathcal{P}}(y \succ x \succ z) = \emptyset$ and $W \subseteq \mathcal{N}^{\mathcal{P}}(y \succ z \succ x) = \emptyset$. For the rest of sets it holds that $A \cup V = \mathcal{N}$. This is the same situation where (x, y) is trivial.

Case 2: $\{x, y, z\}$ does not have a trivial pair. The cannot-be-medium property of y means that there is not a preference with $x \succ y \succ z$ or $z \succ y \succ x$ in Ω . Then it can be said that there exist preferences such that $x \succ z \succ y$, $y \succ x \succ z$, $z \succ x \succ y$, and $y \succ z \succ x$. That is because an isolated triple emerges if any one of them is removed. Let $x \succ z \succ y$ be removed for instance, and $y \succ x \succ z$ becomes an isolated triple. If two are removed then we see either an isolated triple or a trivial pair. A trivial pair is also observed when we remove three of these preferences. Thus (4.5) holds for any partition (A, B, V, W) of \mathcal{N} . Take an arbitrary partition (A, B, V, W) , and let $V' = A \cup V$, $W' = B \cup W$, $A' = B' = \emptyset$. We see that (A', B', V', W') is a partition of \mathcal{N} . Then d satisfies the following:

$$2 \geq d_{A' \cup V'}(x, y) + d_{B' \cup W'}(y, z) + d_{V' \cup W'}(z, x) = d_{A \cup V}(x, y) + d_{B \cup W}(y, z) + 1.$$

We round $d_{A \cup V}(x, y)$ and $d_{B \cup W}(y, z)$ in this inequality, and obtain

$$d'_{A \cup V}(x, y) + d'_{B \cup W}(y, z) \leq 1$$

since player 1 is either in $A \cup V$ or in $B \cup W$. This inequality implies that the “equal to or less than 2” part of (4.5) is satisfied at d' . With regard to the “equal to or more than 1” part of (4.5), let $A' = A \cup V$, $V' = B \cup W$, $B' = W' = \emptyset$, and we have a partition (A', B', V', W') of the set of the players. Then d satisfies the following:

$$1 \leq d_{A' \cup V'}(x, y) + d_{B' \cup W'}(y, z) + d_{V' \cup W'}(z, x) = d_{A \cup V}(x, y) + d_{B \cup W}(y, z).$$

We round the d in this inequality and obtain

$$d'_{A \cup V}(x, y) + d'_{B \cup W}(y, z) \geq 1.$$

This observation tells us that d' satisfies the “equal to or more than 1” part of (4.5).

We have confirmed that for a given $d \in \mathcal{O}^{\mathcal{P}}$, the rounded vector d' is also in $\mathcal{O}^{\mathcal{P}}$. Then assuming that there is a noninteger extreme point leads to a contradiction as we have seen in the single-peaked case. \square

5. PROBABILISTIC ASWFs AND LINEAR INEQUALITY REPRESENTATION

In this section we discuss the relation between a noninteger solution to the linear inequality representation and a probabilistic social welfare function. A probability measure on $L(\mathcal{X})$ is a function $\ell : 2^{L(\mathcal{X})} \rightarrow [0, 1]$ such that $\ell(\emptyset) = 0$, $\ell(L(\mathcal{X})) = 1$, and $\ell(\omega_1 \cup \omega_2) = \ell(\omega_1) + \ell(\omega_2) - \ell(\omega_1 \cap \omega_2)$ for all $\omega_1, \omega_2 \in 2^{L(\mathcal{X})}$. We let $\mathcal{L}(L(\mathcal{X}))$ be the set of all probability measures on $L(\mathcal{X})$.

Definition 5.1. A probabilistic social welfare function (on linear orderings) is a mapping, say h , that maps each profile to a probability measure on $L(\mathcal{X})$, that is, $h : \mathcal{P} \rightarrow \mathcal{L}(L(\mathcal{X}))$, where $\mathcal{P} = \Omega^n \subseteq L(\mathcal{X})^n$. The probabilistic social welfare function h is said to satisfy *unrestricted domain property* when $\Omega = L(\mathcal{X})$.

For each profile $p \in \mathcal{P}$ and for each pair of distinct alternatives $x, y \in \mathcal{X}$, we define $r^h(p, x \succ y)$ as the sum of probabilities that h gives to the preference orderings such that $x \succ y$ at profile p , i.e.,

$$r^h(p, x \succ y) := \ell(\{\triangleright \in L(\mathcal{X}) \mid \triangleright|_{\{x,y\}} = x \succ y\}),$$

where $\ell = h(p) \in \mathcal{L}(L(\mathcal{X}))$. We call the function h satisfying the probabilistic version of Arrow's two axioms shown below a *probabilistic Arrovian Social Welfare Function* (probabilistic ASWF for short).

Axiom 5.2 (Pareto Principle (PP)). If the property that

$$x \succ_i^p y \text{ for all } i \in \mathcal{N} \text{ implies } r^h(p, x \succ y) = 1$$

holds for any pair of distinct alternatives $x, y \in \mathcal{X}$ and for any profile $p \in \mathcal{P}$, then the probabilistic social welfare function h is said to have the *Pareto principle*.

Axiom 5.3 (Independence of Irrelevant Alternatives (IIA)). If the property that

$$\succ_i^{p_1}|_{\{x,y\}} = \succ_i^{p_2}|_{\{x,y\}} \text{ for all } i \in \mathcal{N} \text{ implies } r^h(p_1, x \succ y) = r^h(p_2, x \succ y)$$

holds for any pair of distinct alternatives $x, y \in \mathcal{X}$ and for any pair of distinct profiles $p_1, p_2 \in \mathcal{P}$, then the probabilistic social welfare function h is said to satisfy *independence of irrelevant alternatives*.

We show that a given probabilistic ASWF induces a real vector satisfying the inequalities of the ASWF representation and it coincides with a subadditive function of Barberá and Sonnenschein [4] on the family of subsets of players under unrestricted domain property. The way of constructing the vector is as this: given an $h : \mathcal{P} \rightarrow \mathcal{L}(L(\mathcal{X}))$, we set

$$(5.1) \quad d_S(x, y) := \frac{\sum_{p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)} r^h(p, x \succ y)}{|\mathcal{C}^{\mathcal{P}}(S, x \succ y)|}$$

for each pair of distinct alternatives $x, y \in \mathcal{X}$ and for each set S such that $\mathcal{N}^{\mathcal{P}}(x \succ y) \setminus \mathcal{N}^{\mathcal{P}}(y \succ x) \subseteq S \subseteq \mathcal{N}^{\mathcal{P}}(x \succ y)$.

Theorem 5.4. *For every probabilistic ASWF, $d_S(x, y)$ defined by (5.1) is a nonnegative solution to (3.4)-(3.6).*

Proof. From the definition of r^h

$$r^h(p, x \succ y) + r^h(p, y \succ x) = 1$$

and

$$\begin{aligned} r^h(p, x \succ y) + r^h(p, y \succ z) &= r^h(p, x \succ y \text{ or } y \succ z) + r^h(p, x \succ y \text{ and } y \succ z) \\ &\leq 1 + r^h(p, x \succ z) \end{aligned}$$

hold for any profile $p \in \mathcal{P}$ and for any distinct $x, y, z \in \mathcal{X}$. We simply let

$$d(p, x, y) := r^h(p, x \succ y),$$

then (3.11) as well as (3.12) in Section 3 is satisfied. We also see that (3.9) holds because of (PP) of Axiom 5.2, and (3.10) holds because of (IIA) of Axiom 5.3. As we have seen in the proof of Lemma 3.5 that $d_S(x, y)$ defined by $d(p, x, y)$ through (3.13) satisfies (3.4)-(3.6), $d_S(x, y)$ defined by (5.1) satisfies (3.4)-(3.6). \square

Note that under (IIA) of Axiom 5.3, $d_S(x, y)$ of (5.1) satisfies

$$(5.2) \quad d_S(x, y) = r^h(p, x \succ y) \quad \text{for any } p \in \mathcal{C}^{\mathcal{P}}(S, x \succ y)$$

or equivalently

$$(5.3) \quad d_{\mathcal{S}(p, x \succ y)}(x, y) = r^h(p, x \succ y).$$

While the following statement can be found in Sethuraman et al. [11], we give the proof to make this paper self-contained.

Lemma 5.5. *When the domain is unrestricted, the solution $d_S(x, y)$ to (3.4)-(3.6) does not depend on the pair (x, y) of alternatives.*

Proof. We consider (3.6) for an arbitrary triple of distinct alternatives $x, y, z \in \mathcal{X}$. By unrestricted domain property, $\mathcal{N}^{\mathcal{P}}(x \succ z \succ y)$, $\mathcal{N}^{\mathcal{P}}(y \succ x \succ z)$, $\mathcal{N}^{\mathcal{P}}(z \succ y \succ x)$, $\mathcal{N}^{\mathcal{P}}(x \succ y \succ z)$, $\mathcal{N}^{\mathcal{P}}(z \succ x \succ y)$, and $\mathcal{N}^{\mathcal{P}}(y \succ z \succ x)$ are all \mathcal{N} . Take an arbitrary $S \subseteq \mathcal{N}$ and let $U = S$, $W = \mathcal{N} \setminus S$, the other sets be empty. Then the following holds:

$$d_S(x, y) + d_{S \cup (\mathcal{N} \setminus S)}(y, z) + d_{\mathcal{N} \setminus S}(z, x) \leq 2.$$

Applying (3.4) and (3.5) to this inequality ensures

$$d_S(x, y) \leq d_S(x, z).$$

Meanwhile, we set $U = S$, $V = \mathcal{N} \setminus S$ and have

$$d_S(y, z) \leq d_S(x, z).$$

Since x, y , and z are arbitrarily chosen, these consequences say that the value of $d_S(x, y)$ does not depend on the pair of alternatives but only the set S . \square

Thus, when the domain is unrestricted, the system (3.4)-(3.6) reduces to

$$(5.4) \quad d_{\mathcal{N}} = 1,$$

$$(5.5) \quad d_S + d_{\mathcal{N} \setminus S} = 1 \quad \text{for each } S \subseteq \mathcal{N},$$

$$(5.6) \quad d_{A \cup U \cup V} + d_{B \cup U \cup W} + d_{C \cup V \cup W} \leq 2 \quad \text{for each partition } (A, B, C, U, V, W) \text{ of } \mathcal{N}.$$

Barberá and Sonnenschein [4] characterized the probabilistic ASWF as follows.

Theorem 5.6 (Barberá and Sonnenschein [4]). *For every probabilistic ASWF h on the unrestricted domain, there exists a function $\mu : 2^{\mathcal{N}} \rightarrow \mathbb{R}$ such that*

$$\mu(\mathcal{S}(p, x \succ y)) = r^h(p, x \succ y)$$

for each profile $p \in \mathcal{P}$ and for each pair of distinct alternatives $x, y \in \mathcal{X}$, and furthermore μ satisfies

$$(5.7) \quad \mu(\mathcal{N}) = 1,$$

$$(5.8) \quad \mu(S) + \mu(\mathcal{N} \setminus S) = 1 \quad \text{for each } S \subseteq \mathcal{N},$$

$$(5.9) \quad \mu(S \cup T) \leq \mu(S) + \mu(T) \quad \text{for each } S, T \subseteq \mathcal{N} \text{ (subadditivity)}.$$

Since we have seen (5.3), to see this theorem we have only to show the equivalence between the system (5.4)-(5.6) and (5.7)-(5.9).

Lemma 5.7. *When the domain is unrestricted, the systems (5.4)-(5.6) and (5.7)-(5.9) are equivalent.*

Proof. Suppose d satisfies (5.4)-(5.6) and let S and T be arbitrary subsets of \mathcal{N} . We set $C = S \cap T, U = \mathcal{N} \setminus (S \cup T), V = T \setminus S$ and $W = S \setminus T$. Then from (5.4)-(5.6) we see

$$d_{S \cup T} \leq d_S + d_T.$$

Next suppose μ satisfies (5.7)-(5.9) and let a partition (A, B, C, U, V, W) be given. Then

$$\begin{aligned} \mu(A \cup U \cup V) + \mu(B \cup U \cup W) + \mu(C \cup V \cup W) \\ &= 1 - \mu(B \cup C \cup W) + 1 - \mu(A \cup C \cup V) + 1 - \mu(A \cup B \cup U) \\ &\leq 3 - \mu(A \cup B \cup C \cup U \cup V \cup W) \\ &= 2, \end{aligned}$$

which is (5.6). □

Note that the following monotonicity condition in the original version of Theorem 5.6:

$$\mu(S') \leq \mu(S) \text{ for each } S, S' \subseteq \mathcal{N} \text{ with } S' \subseteq S,$$

can be derived from (5.7)-(5.9) by setting $T = \mathcal{N} \setminus S'$ as follows:

$$\mu(S) + \mu(\mathcal{N} \setminus S') \geq \mu(S \cup (\mathcal{N} \setminus S')) = \mu(\mathcal{N}) = 1 = \mu(S') + \mu(\mathcal{N} \setminus S').$$

We next consider the construction of a probabilistic ASWF from a solution to the linear inequalities. Let $\mathcal{H}^{\mathcal{P}}$ be the set of all probabilistic ASWFs on the given $\mathcal{P} = \Omega^n \subseteq L(\mathcal{X})^n$, and recall that $\mathcal{O}^{\mathcal{P}}$ defined in the proof of Theorem 4.9 is the set of nonnegative solutions satisfying (3.4)-(3.6). We denote by $\Gamma^{\mathcal{P}} : \mathcal{H}^{\mathcal{P}} \rightarrow \mathcal{O}^{\mathcal{P}}$ a function that assigns a nonnegative solution in $\mathcal{O}^{\mathcal{P}}$ to a probabilistic ASWF by (5.1). Concerning its characteristics, the first question is whether $\Gamma^{\mathcal{P}}$ is injective. The following example shows that it is not injective in general.

Example 5.8. Let $n = 2$, $\mathcal{X} = \{x, y, z\}$, and $\Omega = \{x \succ y \succ z, z \succ y \succ x\}$. Then there are four profiles, named p_1, \dots, p_4 . We define two probabilistic ASWFs h_1, h_2 where the probability of each social preference being selected at each profile is shown below. While these two functions h_1 and h_2 are different, they give the same value of r^h and hence $d_S(x, y)$. In fact, take p_2 for example, then we have

$$r^{h_1}(p_2, x \succ y) = h_1(p_2) (\{x \succ y \succ z, x \succ z \succ y, z \succ x \succ y\}) = \frac{1}{2} + 0 + 0 = \frac{1}{2},$$

and

$$r^{h_2}(p_2, x \succ y) = h_2(p_2) (\{x \succ y \succ z, x \succ z \succ y, z \succ x \succ y\}) = \frac{1}{4} + \frac{1}{4} + 0 = \frac{1}{2}.$$

$h(p)$		$x \succ y \succ z$	$z \succ y \succ x$	$x \succ z \succ y$	$y \succ z \succ x$	$y \succ x \succ z$	$z \succ x \succ y$
h_1	$x \succ_1^{p_1} y \succ_1^{p_1} z$	1	0	0	0	0	0
	$x \succ_2^{p_1} y \succ_2^{p_1} z$						
h_1	$x \succ_1^{p_2} y \succ_1^{p_2} z$	1/2	1/2	0	0	0	0
	$z \succ_2^{p_2} y \succ_2^{p_2} x$						
h_1	$z \succ_1^{p_3} y \succ_1^{p_3} x$	1/2	1/2	0	0	0	0
	$x \succ_2^{p_3} y \succ_2^{p_3} z$						
h_1	$z \succ_1^{p_4} y \succ_1^{p_4} x$	0	1	0	0	0	0
	$z \succ_2^{p_4} y \succ_2^{p_4} x$						
h_2	$x \succ_1^{p_1} y \succ_1^{p_1} z$	1	0	0	0	0	0
	$x \succ_2^{p_1} y \succ_2^{p_1} z$						
h_2	$x \succ_1^{p_2} y \succ_1^{p_2} z$	1/4	1/4	1/4	1/4	0	0
	$z \succ_2^{p_2} y \succ_2^{p_2} x$						
h_2	$z \succ_1^{p_3} y \succ_1^{p_3} x$	1/4	1/4	1/4	1/4	0	0
	$x \succ_2^{p_3} y \succ_2^{p_3} z$						
h_2	$z \succ_1^{p_4} y \succ_1^{p_4} x$	0	1	0	0	0	0
	$z \succ_2^{p_4} y \succ_2^{p_4} x$						

TABLE 1. Two different probabilistic ASWFs that yield the same $d_S(x, y)$

The second question is whether $\Gamma^{\mathcal{P}}$ is surjective, i.e., if there is a probabilistic ASWF for each point of $\mathcal{O}^{\mathcal{P}}$. Concerning the problem, we should observe the result of McLennan [7] that when the number of alternatives is six or more and the domain is unrestricted, the function μ in Theorem 5.6 is additive, that is,

$$(5.10) \quad \mu(S \cup T) = \mu(S) + \mu(T) \text{ for all disjoint } S, T \subseteq \mathcal{N}.$$

Using this result we see by the next example that $\Gamma^{\mathcal{P}}$ is not surjective in general.

Example 5.9. Let $n \geq 3$, $|\mathcal{X}| \geq 6$, and $\mathcal{P} = \Omega^n = L(\mathcal{X})^n$, i.e., unrestricted domain. For each pair of distinct alternatives $x, y \in \mathcal{X}$, and for each set $S \subseteq \mathcal{N}$, we set

$$d_S(x, y) := \begin{cases} 1 & \text{if } S = \mathcal{N}, \\ \frac{1}{2} & \text{if } \emptyset \subsetneq S \subsetneq \mathcal{N}, \\ 0 & \text{if } S = \emptyset. \end{cases}$$

This is clearly a fractional solution to (3.4)-(3.6). We suppose that this is obtained by a probabilistic ASWF h through (5.1). Then, as we have seen before $d_{S(p, x \succ y)}(x, y) = r^h(p, x \succ y)$, and from Theorem 5.6 we also have $r^h(p, x \succ y) = \mu(\mathcal{S}(p, x \succ y))$. Since the domain is unrestricted, we obtain that for every pair of distinct alternatives x and y and for every $S \subseteq \mathcal{N}$

$$d_S(x, y) = \mu(S).$$

Note that $d_S(x, y)$ is independent of the pair of alternatives. Let \mathcal{N} be partitioned into three nonempty sets, say S_1, S_2, S_3 , then $d_{S_1} + d_{S_2} = 1/2 + 1/2 = 1$ whereas $d_{S_1 \cup S_2} = d_{\mathcal{N} \setminus S_3} = 1/2$. We see that $\mu(S) = d_S$ does not satisfy (5.10) and hence not an image of $\Gamma^{\mathcal{P}}$.

The following result shows that we have only to check the extreme points of the polytope when we judge whether $\Gamma^{\mathcal{P}}$ is surjective.

Theorem 5.10. *The function $\Gamma^{\mathcal{P}}$ is surjective if and only if for each extreme point d^* of $\mathcal{O}^{\mathcal{P}}$ there exists a probabilistic ASWF h^* such that $\Gamma^{\mathcal{P}}(h^*) = d^*$.*

Proof. The “only if” part is trivial, and we show the “if” part. For a given \mathcal{P} we denote all the extreme points of polytope $\mathcal{O}^{\mathcal{P}}$ by $d_1^*, \dots, d_{\mathcal{K}}^*$. Then for any point \bar{d} in $\mathcal{O}^{\mathcal{P}}$ there is $\lambda_1, \dots, \lambda_{\mathcal{K}}$ satisfying $\sum_{i=1}^{\mathcal{K}} \lambda_i = 1$ as well as $\lambda_i \geq 0$ for each $i \in \{1, \dots, \mathcal{K}\}$ and \bar{d} can be written as

$$\bar{d} = \sum_{i=1}^{\mathcal{K}} \lambda_i d_i^*.$$

For each $i \in \{1, \dots, \mathcal{K}\}$ let h_i^* a probabilistic ASWF with $\Gamma^{\mathcal{P}}(h_i^*) = d_i^*$. We define \bar{h} as

$$\bar{h}(p)(\triangleright) := \sum_{i=1}^{\mathcal{K}} \lambda_i h_i^*(p)(\triangleright)$$

for each $p \in \mathcal{P}$ and for each $\triangleright \in L(\mathcal{X})$, then \bar{h} is a probabilistic ASWF and $\Gamma^{\mathcal{P}}(\bar{h}) = \bar{d}$. \square

Let us consider the polytope determined by (5.7)-(5.10) and the nonnegativity constraint of μ . Combining (5.7) and (5.10), we have (5.8). Furthermore (5.9) is obtained from (5.10) and nonnegativity:

$$\begin{aligned} \mu(T) + \mu(S) - \mu(S \cup T) &= \mu(T) + \mu(S \cap T) + \mu(S \setminus T) - \mu(S \setminus T) - \mu(T) \\ &= \mu(S \cap T) \geq 0. \end{aligned}$$

Thus (5.7)-(5.10) together with the nonnegativity constraint is equivalent to the system of (5.7), (5.10) and the nonnegativity. Therefore there is an n -dimensional vector $\pi = (\pi_1, \dots, \pi_n)$ in the regular simplex $\Pi := \{ \pi \in \mathbb{R}^n \mid \pi \geq 0; \sum_{i=1}^n \pi_i = 1 \}$ such that

$$(5.11) \quad \mu(S) = \sum_{i \in S} \pi_i$$

holds for every $S \subseteq \mathcal{N}$. Conversely, for a given $\pi \in \Pi$, $\mu(S) := \sum_{i \in S} \pi_i$ satisfies (5.7), (5.10) and nonnegativity. Therefore the polytope defined by (5.7)-(5.10) and the nonnegativity is the image of the regular simplex $\Pi \subseteq \mathbb{R}^n$ under the linear function defined by (5.11). Since each extreme point of the polytope is an image of an extreme point of Π , which is a unit vector, we see that all the extreme points of the polytope are integer vectors. In fact, for each $i \in \mathcal{N}$ let $\mu^i \in \mathbb{R}^{2^n - 1}$ be defined by

$$\mu^i(S) := \begin{cases} 1 & \text{if } i \in S, \\ 0 & \text{otherwise.} \end{cases}$$

Then the set of extreme points of the polytope consists of μ^i 's for $i \in \mathcal{N}$. Since, as we have seen, an integer extreme point of the polytope is an image of an ASWF, this together with Theorem 5.10 means that $\Gamma^{\mathcal{P}}$ is surjective in this case.

6. CONCLUDING REMARKS

This paper presented a study on the linear inequality representation of Arrowian Social Welfare Functions. We gave an alternative proof of the ASWF integer linear inequality representation theorem of Sethuraman et al. [11]. Our technique of proof can be applied to derive another linear inequality formulation of ASWFs for different preference domain frameworks such as the one discussed in Ando, Ohara, and Yamamoto [1] and Ohbo, Tsurutani, Umezawa, and Yamamoto [8] as well as an ASWF on weak orderings. See Sato [9] for further details. We also studied the polyhedral structure determined by the linear inequalities on single-peaked domains, single-caved domains, and the domains where each triple of alternatives contains one that cannot be medium, showing that the set of

nonnegative solutions to the inequalities forms an integral polytope when these domains satisfy weakly nonisolated condition. We then showed that a real vector satisfying the linear inequalities can be created from any probabilistic ASWF and derived the subadditive function of Barberá and Sonnenschein [4] as a special case. We also considered the construction of a probabilistic ASWF from a given nonnegative solution to the inequalities.

There still remain interesting problems unsolved. One is the characterization of social choice functions as a system of inequalities. A construction problem of a probabilistic ASWF from a solution to the linear inequalities on restricted domains is also worth further study. Study of the probabilistic version of social choice functions in Gibbard [5] through inequality representations is left for future research.

ACKNOWLEDGEMENTS

The authors thank René van den Brink and Gerard van der Laan for stimulating discussion. This paper was written while the second author was visiting professor at the Tinbergen Institute of the Faculty of Economics and Business Administration, Vrije Universiteit, Amsterdam, the Netherlands with the grant B 46-551 from the Netherlands Organisation for Scientific Research. They are very grateful to the organisation as well as the university.

REFERENCES

- [1] K. Ando, A. Ohara, and Y. Yamamoto. Impossibility theorems on mutual evaluation. *Journal of the Operations Research Society of Japan*, 46(4):523–532, 2003. (in Japanese).
- [2] K. J. Arrow. *Social Choice and Individual Values*. Wiley, New York, 2nd edition, 1963.
- [3] K. J. Arrow, A. K. Sen, and K. Suzumura, editors. *Handbook of Social Choice and Welfare*, volume 1. Elsevier, Amsterdam, 2002.
- [4] S. Barberá and H. Sonnenschein. Preference aggregation with randomized social orderings. *Journal of Economic Theory*, 18:244–254, 1978.
- [5] A. Gibbard. Manipulation of schemes that mix voting with chance. *Econometrica*, 45(3):665–681, 1977.
- [6] E. Kalai and E. Muller. Characterization of domains admitting nondictatorial social welfare functions and nonmanipulable voting procedures. *Journal of Economic Theory*, 16:457–469, 1977.
- [7] A. McLennan. Randomized preference aggregation: Additivity of power and strategy proofness. *Journal of Economic Theory*, 22:1–11, 1980.
- [8] K. Ohbo, M. Tsurutani, M. Umezawa, and Y. Yamamoto. Social welfare function for restricted individual preference. *Pacific Journal of Optimization*, 1(2):315–325, 2005.
- [9] K. Sato. A study on linear inequality representation of social welfare functions. Master’s thesis, Graduate School of Systems and Information Engineering, University of Tsukuba, January 2006.
- [10] A. K. Sen. *Collective Choice and Social Welfare*. Holden-Day, San Francisco, 1970.
- [11] J. Sethuraman, C. P. Teo, and R. V. Vohra. Integer programming and Arrovian social welfare functions. *Mathematics of Operations Research*, 28(2):309–326, 2003.

KEISUKE SATO AND YOSHITSUGU YAMAMOTO: GRADUATE SCHOOL OF SYSTEMS AND INFORMATION ENGINEERING, UNIVERISTY OF TSUKUBA, TSUKUBA, IBARAKI 305-8573, JAPAN
E-mail address: {sato90, yamamoto}@sk.tsukuba.ac.jp

Tinbergen Institute

The Tinbergen Institute is the institute for economic research of the Erasmus Universiteit Rotterdam, Universiteit van Amsterdam, and Vrije Universiteit Amsterdam.

Tinbergen Institute Amsterdam

Roetersstraat 31

1018 WB Amsterdam

The Netherlands

Tel.: +31(0)20 551 3500

Fax: +31(0)20 551 3555

Tinbergen Institute Rotterdam

Burg. Oudlaan 50

3062 PA Rotterdam

The Netherlands

Tel.: +31(0)10 408 8900

Fax: +31(0)10 408 9031

Please send questions and/or remarks of non-scientific nature to driessen@tinbergen.nl.

Most TI discussion papers can be downloaded at <http://www.tinbergen.nl>.